

# Recitation 1

## *Proofs and Logic*


### What's recitation?

Recitation is a space for you to work with other members of the CS22 community on problems that we hope will help you hone your understanding of the course material, get better at communicating with other folks about mathematical ideas, and practice for the homework. You'll also get to know some of the TAs and ask them any questions about the course material that you're passionate about, barring specific questions about the homework for the week. If you have any feedback on recitation or the course in general, please share with us through the [anonymous feedback form](#).

### Check-in

**Stop!** Read this section in its entirety before beginning recitation!

We will be using *Gradescope* for getting checked off for recitation. Right now, **one** member of your group should log into Gradescope, select the *Recitation 1* assignment, and submit a blank assignment (ignore all inputs):



The screenshot shows a Gradescope question interface. At the top, it says "Q2 Final check-off" with "0.5 Points" below it. The instructions read: "Ignore this and just submit the assignment (leave this blank!). Your grade for this question will update once you get your final check-off." Below the instructions is a text input field with the placeholder "Enter your answer here". There are three buttons: "Save Answer" below the input field, "Save All Answers" at the bottom left, and "Submit & View Submission >" at the bottom right. A blue arrow points from the text "Click Here!" to the "Submit & View Submission >" button, which is also highlighted with a blue rounded rectangle.

Then, in the next screen, select that you want to add group members:

Finally, add your group members into your submission:

This will mark that you're here at recitation with your group! You're all set!

# Proof Techniques

## Direct Proof

A direct proof is one that begins with statements you know to be true, makes logical jumps to new statements, and eventually ends up with what you are trying to prove.

Here is an example:

### Example

**Claim:** If  $n$  is odd, then  $n^2$  is odd.

*Proof.* We know that  $n$  is odd, so  $n = 2k + 1$  for some  $k \in \mathbb{Z}$ .

So  $n^2 = (2k + 1)(2k + 1) = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1 = 2m + 1$ , where  $m = 2k^2 + 2k$ .

Since  $m$  is an integer,  $n^2$  is odd. □

a. Prove that the product of an even number and odd number is even.



b. *Optional:* Prove that the product of two rational numbers is rational.

## Negation and Counterexample

Sometimes, we don't ask you to prove claims are true; we instead ask you to prove that they are false! This means your task is to prove the *negation* of the claim.

What is the negation of a claim? Let's think about it.

**Question:** Suppose Pete says to Joe: "Everyone at Brown likes sunflowers!". What would Joe need to show Pete to convince him this was false? Circle your choice.

- (a) Every person at Brown hates sunflowers.
- (b) There is at least one person at Brown who does not like sunflowers.
- (c) There are at least 22 people at Brown who do not like sunflowers.

**Question:** Now, practice negating each of the following claims. Write the negated version of the claim below.

- a. All CS22 students want to be a gardener.
- b. There exists a student in CS22 who is a gardener.
- c.  $\forall x \in \mathbb{Z}$ , if  $x$  is even,  $2x$  is odd.

Sometimes, we can show that a claim is false (i.e. the negation of the claim is true) by providing a **counterexample**.

### Example

Suppose Joe makes the claim that, if  $xy$  is rational then  $x$  and  $y$  are rational.

Pete can disprove Joe's claim by coming up with a counterexample. He notices that Joe's statement has an implicit "for all" quantifier: "For all  $x$  and  $y$ , if  $xy$  is rational, then  $x$  and  $y$  are rational." Pete can then negate this statement to get "There exists  $x$  and  $y$  such that  $xy$  is rational and  $x$  or  $y$  is irrational." If Pete chooses  $x = \sqrt{2}$  and  $y = \sqrt{2}$ , then  $x$  and  $y$  are irrational, and  $xy = 2$ , which is rational. This proves the negated claim, and thus disproves Joe's original claim. Pete's counterexample was  $x = \sqrt{2}$  and  $y = \sqrt{2}$ .

**Question:** Now, disprove the following statement by providing a counterexample. If  $xyz$  is rational, then  $x$ ,  $y$ , and  $z$  are rational.

**Optional Recommended Checkpoint** — Call over a TA.

## Logic

1. A **propositional formula** is a function of one or more variables, each of which can be set to true or false, that evaluates to true or false. We call a propositional formula a *proposition* for short.
2. The term **logical expression** is often used synonymously with the word proposition.
3. Two propositions are **logically equivalent** when they have the same truth tables.
4. A proposition is **valid** if it evaluates to true on any choice of inputs; it is true no matter what. That is, a valid proposition is logically equivalent to the expression  $(p \text{ OR NOT } p)$ . This is also called a *tautology*.
5. A proposition is **satisfiable** if it evaluates to true on some choice of inputs. A valid proposition is satisfiable, but so are many propositions which sometimes evaluate to false.
6. If a proposition is not satisfiable, it evaluates to false on any choice of inputs; it is false no matter what. That is, it is logically equivalent to the expression  $(p \text{ AND NOT } p)$ . This is called a *contradiction*.

Let's now review the interpretation of each of the following logical connectives:

$p$	$q$	$\neg p$	$p \wedge q$	$p \vee q$	$p \text{ xor } q$	$p \rightarrow q$	$p \leftrightarrow q$
T	T	F	T	T	F	T	T
T	F	F	F	T	T	F	F
F	T	T	F	T	T	T	F
F	F	T	F	F	F	T	T

### Why do we have propositions?

Propositions are condensed ways of representing truth tables. Because they're more condensed, we oftentimes can more easily extract the meaning from them. It can be hard to look at a truth table and see the "general rule" that transforms input to output. A proposition is that rule.

Remember when we represented functions in the form  $f(x) = \dots$ ? This notation is a general rule for creating the set of ordered pairs that is the function. In the same way, a proposition is a general rule for creating the truth table.

## Logical Equivalence: Two approaches

You have two techniques at your disposal to determine if two expressions are logically equivalent: by using truth tables or by using logical rewrite rules.

Given two expressions, you can write out the truth table for each one. If they have the same inputs and their truth tables are the same, they are logically equivalent.

Or, given two expressions, you use logical equivalence rules to try to get from one expression to the other. For example, take the expression  $\neg(x \wedge y)$ . Using DeMorgan's law, you could get that the not “distributes” in this expression, and it is therefore equivalent to  $(\neg x \vee \neg y)$ . A full list of rules you can use can be found [on the course website](#). Read the  $\equiv$  symbol as “is equivalent to.”

**Task:** Determine if each pair of expressions below are logically equivalent (i.e. determine if the equivalence stated holds). You can choose to either use truth tables or simplify the expressions to arrive at your answer. To show that the expressions are not logically equivalent, you only need to provide one counterexample.

a.  $p \vee q \equiv (\neg p) \wedge (\neg q)$

b.  $p \leftrightarrow q \equiv (p \wedge q) \vee (\neg p \wedge \neg q)$



c. *Optional:*  $p \wedge \neg q \equiv \neg(\neg p \vee q)$



d. *Optional:*  $((p \vee q) \wedge z) \equiv (p \vee (q \wedge z))$

## More Practice

- a. Give an assignment to the variables  $x_1, x_2, x_3$  which makes the following logical expression evaluate to true.

$$(x_1 \vee x_2) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3) \wedge x_3$$

- b. Is the following logical expression *valid*? That is, is it true under every truth assignment? Explain your answer.

$$(x_1 \wedge x_2) \vee (\neg x_1 \wedge \neg x_2 \wedge \neg x_3) \vee (x_3 \wedge \neg x_3)$$

- c. Come up with a logical expression with three variables which has **only one** assignment to the variables which makes it true.

# Facebook Censorship Rules

In 2017, ProPublica allegedly released information on how Facebook determines what constitutes hate speech. The ruleset they outlined was a highly deterministic, logic-based system (note the difference between logic-based and logical).

Here are several overarching censorship rules that Facebook follows to determine if a comment is hate speech (note that enforcement is inconsistent at best, and it is unknown whether or not Facebook has changed its methodology):

1. A comment in question is making an “attack” or hateful comment (the specifics on how this is determined is not clear).
2. The comment in question is directed towards **people** belonging to a “protected group” or any combination of protected groups. Note that each member of a protected group is protected indiscriminately (comments disparaging “man” are weighed equally as comments disparaging “woman”) and the comment must be directed towards people, not ideologies.
  - The protected groups are race, sex, gender identity, religious affiliation, national origin, ethnicity, sexual orientation, and serious disability/disease. This group can shift depending on current events; for example, “Syrian immigrants” were added as a protected group in the wake of the Syrian refugee crisis.
3. If the target of the comment has multiple listed characteristics, (e.g. “Asian” has one characteristic, “Asian male” has two, and “elderly Asian male” has three), **each characteristic must be in a protected group** for the comment to be considered hate speech.

If these conditions are met, Facebook flags the comment as hate speech and the comment is subject to being taken down. We can express this as a proposition: let  $p$  be that the comment is hateful, let  $q_i$  be that the  $i$ th characteristic of the targeted group belongs in the “protected groups,” and let  $r$  be that the given comment is hate speech. We apply this simple logic to determine whether the statement is hate:

$$p \wedge q_1 \wedge q_2 \wedge q_3 \cdots \rightarrow r$$

**Task:** Given the following list of subgroup characteristics, determine if the following groups would be protected from “hate speech” under the Facebook rules (focus on rules three and two). Give another example of a group that should be protected but isn’t.

1. Female driver
2. Black kid
3. White man
4. Islam
5. Muslims

**Task:** In 2018, Facebook responded to the article by adding “age” into the list of protected groups. How have your answers to the previous tasks changed? How could someone continue to post racist, sexist, and hateful comments with these rules? Does something still strike you wrong about the list of protected groups and unprotected groups?

**Task:** Facebook tried its best to use a rigid set of predicates and propositions to determine whether a given comment is hate speech. Describe one limitation that logical formulas have in terms of applying them to complicated real-world situations.



## Proof by Contradiction

In math, a claim and its negation cannot both be true. This would be a **contradiction**. For example, it cannot be the case that 3 is odd and 3 is not odd. Only one of these statements can be true, and one of them has to be true.

This idea motivates a proof technique called *proof by contradiction*. The general idea is this:

Say we have some statement  $T$  that we are trying to prove is true.

To prove  $T$  is true by contradiction:

1. We begin by assuming  $T$  is NOT true. That is, we assume that the negation of  $T$  is true.
2. Assuming  $T$  is not true leads us to a contradiction. That is, by making logical leaps from  $T$  being not true, we arrive at the fact that a statement  $x$  and its negation both have to be true.
3. But  $x$  and its negation cannot both be true; this is a contradiction. We got to this contradiction by assuming  $T$  was false. Therefore, we know  $T$  cannot be false; i.e.  $T$  is true.

Here is an example.

### Example

**Claim:** There is no smallest rational number greater than 0.

*Proof.* Assume for sake of contradiction that there exists a smallest rational number, say  $r$ .

However,  $r/2$  is a rational number greater than 0 and smaller than  $r$ .

This is a contradiction to the fact that  $r$  is the smallest rational number.

Assuming that there is a smallest rational number led to a contradiction, and therefore there is no smallest rational number greater than 0.  $\square$

Now it's your turn! Prove the following statements by contradiction.

- a. 2 is an even number.
- b. *Optional:* Suppose  $a, b \in \mathbb{R}$ . If  $a$  is rational and  $ab$  is irrational, then  $b$  is irrational.



## Proof by Contrapositive

If we have a statement “if  $A$  then  $B$ ”, we define the contrapositive of that statement to be “if not  $B$  then not  $A$ ”. If the statement is true, then the contrapositive is true, and if the statement is false, then the contrapositive is false (check this for yourself!). We can therefore prove statements just by showing that their contrapositive is true.

### Example

Let’s say we want to prove the statement *if  $x^2$  is even, then  $x$  is even*. We could spend the effort to come up with a new direct proof of this. But, earlier in the recitation (the direct proof example) we already showed that *if  $n$  is odd, then  $n^2$  is odd*. This is exactly the contrapositive of what we want to show, so we can just reuse that proof for this new statement.


**Question:** Now, practice writing the contrapositive for each of these statements. Write the contrapositive next to the original.

1. If a student attends recitation, then they understand the material.
2. If someone is a gardener, then they grow tulips.
3. A student is a Gemini if they are born on May 21.

**Checkpoint — Call a TA over.**

**Checkoff - call over a TA!**

If you were checked off, you should be able to see this reflected on Gradescope:

NAME	STATUS	RELEASED	DUE (EST)
Recitation 1	1.0 / 1.0	 FEB 03	1 week, 2 days left FEB 14 AT 12:00AM

You will do this for every recitation going forward.

If you weren’t able to complete recitation, you may come to *any* recitation in the next week to get it checked off (optimally with your group). We *highly recommend* you to do this at the same time next week, with your group, *after* you have finished next week’s recitation. The deadline for recitation checkoff is by your next scheduled recitation time.